



CME POSTTEST

EXPIRATION DATE: JUNE 2010

All posttests must be submitted online to AAPA.

The CME scoring service is provided at no charge to current Fellow members of AAPA. Others must submit a \$20 scoring fee for each posttest using AAPA's secure server. Payment may be made using Visa, MasterCard, or American Express. To submit this posttest and earn 1 hour of AAPA Category I CME credit, go to www.aapa.org/cme/post-test.html.

KNEE EXAMINATION (PAGE 20)

1. The valgus stress test is used to test the
 - a. Medial collateral ligament (MCL)
 - b. Anterior cruciate ligament (ACL)
 - c. Posterior meniscofemoral ligament
 - d. Lateral collateral ligament
2. The most accurate clinical sign of a torn meniscus is
 - a. Knee effusion
 - b. Pain with forced hyperextension
 - c. Joint-line tenderness
 - d. History of catching or locking
3. The Ege's test is a weight-bearing version of the
 - a. Apley compression test
 - b. Lachman's test
 - c. Anterior drawer test
 - d. McMurray's test
4. A positive finding with the pivot shift test is pathognomonic of
 - a. Posterior cruciate ligament rupture
 - b. ACL insufficiency
 - c. Patella fracture
 - d. MCL tear
5. The gold standard test for diagnosing tears in the posterior cruciate ligament is the
 - a. Posterior drawer test
 - b. Reverse pivot shift
 - c. Posterior sag test
 - d. Patella grind test
6. During a neurovascular examination, knee flexion is used to evaluate the
 - a. Femoral nerve
 - b. Tibial nerve
 - c. Sciatic nerve
 - d. Peroneal nerve

FIFTH DISEASE (PAGE 42)

7. The transmission rate of parvovirus B19 (B19) is about 50% for
 - a. Day-care workers
 - b. Susceptible teachers
 - c. Household contacts
 - d. Health care workers
8. Classic childhood B19 infection manifests as
 - a. A maculopapular and erythematous rash
 - b. Erythematous plaques on both cheeks
 - c. A fine, red and rough-textured rash
 - d. Vesicles with erythema on the palms and soles
9. After B19 infection, a transient drop occurs in
 - a. Platelet count
 - b. Monocyte count
 - c. Absolute neutrophil count
 - d. Reticulocyte count
10. The typical presentation of fifth disease is the acute onset of arthritis, primarily affecting the
 - a. Proximal interphalangeal joint
 - b. Temporomandibular joint
 - c. Acromioclavicular joint
 - d. Femoral acetabular joint
11. B19 infection should be included in the differential of
 - a. Pneumonitis
 - b. Nephrolithiasis
 - c. Acute myocarditis
 - d. Ulcerative colitis
12. The B19 etiology of unexplained fulminant hepatitis was shown when the B19 genome was isolated in 4 of 11 children who had
 - a. Jaundice
 - b. High AST activity
 - c. Low ALT activity
 - d. A high bilirubin level

Successful completion of the self-assessment is required to earn Category I CME credit. Successful completion is defined as a cumulative score of at least 70% correct.