



CME POSTTEST

EXPIRATION DATE: JULY 2009

All posttests must be submitted online to AAPA.

The CME scoring service is provided at no charge to current Fellow members of AAPA. Others must submit a \$20 scoring fee for each posttest using AAPA's secure server. Payment may be made using Visa, MasterCard, or American Express. To submit this posttest and earn 1 hour of AAPA Category I CME credit, go to www.aapa.org/cme/post-test.html.

GENETIC SUSCEPTIBILITY TO VTE, PAGE 20

1. A type of anemia that is a risk factor for venous thromboembolism (VTE) is
 - a. Iron deficiency
 - b. Sickle cell
 - c. β -thalassemia
 - d. Sideroblastic
2. A diagnosis of VTE is confirmed by
 - a. Doppler ultrasonography
 - b. D-dimer concentration
 - c. Venography
 - d. MRI
3. Hyperhomocysteinemia is associated with the use of
 - a. Alpha-blockers
 - b. Statins
 - c. Antifolates
 - d. Benzodiazepines
4. The prevalence of factor V Leiden is highest among
 - a. Hispanics
 - b. US Whites
 - c. Asians
 - d. African-Americans
5. An antithrombin deficiency is suggested by a lack of response to
 - a. Clopidogrel
 - b. Warfarin
 - c. Vitamin K
 - d. Heparin
6. The presence of lupus anticoagulant may cause prolongation of
 - a. Partial thromboplastin time
 - b. Prothrombin time
 - c. Thrombin clotting time
 - d. Bleeding time

LEAD POISONING PREVENTION, PAGE 40

7. Lead-based paint was banned beginning in
 - a. 1973
 - b. 1978
 - c. 1983
 - d. 1988
8. Currently, average blood lead levels (BLL) range from
 - a. 0.5-1 $\mu\text{g}/\text{dL}$
 - b. 1-2 $\mu\text{g}/\text{dL}$
 - c. 2-4 $\mu\text{g}/\text{dL}$
 - d. 4-6 $\mu\text{g}/\text{dL}$
9. Among children aged 1 to 5 years, the highest prevalence of elevated BLL is in
 - a. non-Hispanic blacks
 - b. Hispanics
 - c. Asians
 - d. Whites
10. Exposure to environmental lead is a risk factor in children for
 - a. Autism
 - b. Depression
 - c. Anorexia
 - d. Attention-deficit/hyperactivity disorder
11. Which of the following remains intact despite chronic lead exposure?
 - a. Muscle tone
 - b. Tendon reflexes
 - c. Sensory function
 - d. Level of consciousness
12. Children with a BLL higher than 45 $\mu\text{g}/\text{dL}$ should be treated with
 - a. Dimercaprol
 - b. Succimer
 - c. Nitritotriacetic acid
 - d. Desferoxamine

Successful completion of the self-assessment is required to earn Category I CME credit. Successful completion is defined as a cumulative score of at least 70% correct.